





Comirnaty® (Pfizer/BioNTech) vaccine Summary Sheet

please refer to www.immunisation.ie for full guidance



**COVID-19
VACCINE**
Public Health
Advice



Storage

- -90°C to -60°C undiluted until expiry date
- +2°C to +8°C up to 1 month (“use before” date and time)
- Must be diluted within 2 hours of removal from +2°C to +8°C fridge
- 6 hours once diluted (“discard” date and time)



Comirnaty®

- 0.3ml dose
- 2 dose schedule*
- 21-28 days interval
- People aged 12 and over

DO NOT SHAKE

Preparation and Dilution

Preparation for dilution



- Remove vaccine from fridge
- Check “use before” date
- Inspect vial
- Gently invert 10 times



Dilute vaccine



- Dilute vial with 1.8mL Sodium Chloride 0.9% Solution for Injection
- Slowly remove 1.8mL of air from vial to equalise pressure
- Gently invert the vial 10 times

Label the vial



- Label the vial with “discard time” (6 hours after time of dilution)

Administration

- Check within “discard time”
- Clean top of vial with alcohol swab and allow to dry
- Withdraw dose 0.3ml
- Gently remove air bubbles
- Administer dose intramuscularly



Version 6.0 8 September 2021
Created by: HSE National Immunisation Office

*Please see Immunisation Guidelines for Ireland for specific recommendations <https://www.hse.ie/eng/health/immunisation/hcpinfo/guidelines/covid19.pdf>