

Vinorelbine and CISplatin Therapy- 28 day

INDICATIONS FOR USE:

INDICATION	ICD10	Regimen Code	Reimbursement Status
Adjuvant treatment of patients with completely resected stage IB, II or IIIA non small cell lung cancer (NSCLC)	C34	00343a	Hospital
Treatment of locally advanced recurrent or metastatic NSCLC	C34	00343b	Hospital

TREATMENT:

The starting dose of the drugs detailed below may be adjusted downward by the prescribing clinician, using their independent medical judgement, to consider each patients individual clinical circumstances.

CISplatin is administered on day 1 and day 8 and vinorelbine weekly on day 1, 8, 15 and 22 of a 28 day cycle for 4 cycles or until disease progression or unacceptable toxicity develops.

Facilities to treat anaphylaxis MUST be present when the chemotherapy is administered.

Admin. Order	Day	Drug	Dose	Route	Diluent & Rate	Cycle
1	1,8, 15 and 22	^a Vinorelbine	25mg/m ²	IV infusion	50ml 0.9% NaCl over 15min. Then flush the line with 250ml 0.9% NaCl prior to removing/capping IV access	Every 28 days for 4 cycles
2	1 and 8	^b CISplatin	50mg/m ²	IV infusion	1000ml NaCl 0.9% over 2 hours	Every 28 days for 4 cycles

^aVinorelbine is a neurotoxic chemotherapeutic agent. Refer to NCCP Guidance on the Safe Use of Neurotoxic drugs (including Vinca Alkaloids) in the treatment of cancer.

<https://www.hse.ie/eng/services/list/5/cancer/profinfo/medonc/safetyreview/neurotoxicguidance.pdf>

^bPre and post hydration therapy required for CISplatin

See local hospital policy recommendations.

Suggested prehydration for CISplatin therapy:

- Administer 1000ml NaCl 0.9% over 1 hour
- Administer CISplatin as described above

Post hydration:

- Administer 1000ml NaCl 0.9% with 10mmol magnesium sulphate (MgSO₄) and 20mmol potassium chloride (KCl) over 2 hours

Mannitol 10% may be used to as per local policy to induce diuresis, although there is no conclusive evidence that this is required.

The routine use of furosemide to increase urine flow is not recommended unless there is evidence of fluid overload (3,4).

ELIGIBILITY:

- Indications as above
- Life expectancy > 3 months
- ECOG 0-1

NCCP Regimen: Vinorelbine and CISplatin Therapy -28 day	Published: 20/06/2016 Review: 10/06/2025	Version number: 5
Tumour Group: Lung NCCP Regimen Code: 00343	ISMO Contributor: Prof Maccon Keane	Page 1 of 6

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <http://www.hse.ie/eng/Disclaimer>

This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens

EXCLUSIONS:

- Hypersensitivity to vinorelbine or other vinca alkaloids, cisplatin or any of the excipients
- Moderate/severe renal impairment (creatinine clearance < 60 mL/min)
- Significant hearing impairment/tinnitus
- Pre existing neuropathies \geq grade 2
- Pregnancy
- Lactation

USE with CAUTION:

- Neutrophil count < $1.5 \times 10^9/L$ or severe infection; current or recent (within 2 weeks)
- Platelet count < $100 \times 10^9/L$

PRESCRIPTIVE AUTHORITY:

The treatment plan must be initiated by a Consultant Medical Oncologist

TESTS:

Baseline tests:

- FBC, renal and liver profile
- Audiometry and creatinine clearance as clinically indicated
- Assessment of peripheral neuropathy

Regular tests:

- FBC weekly
- Renal and liver profile prior to each cycle

Disease monitoring:

Disease monitoring should be in line with the patient's treatment plan and any other test/s as directed by the supervising Consultant.

DOSE MODIFICATIONS:

- Any dose modification should be discussed with a Consultant.

NCCP Regimen: Vinorelbine and Cisplatin Therapy -28 day	Published: 20/06/2016 Review: 10/06/2025	Version number: 5
Tumour Group: Lung NCCP Regimen Code: 00343	ISMO Contributor: Prof Maccon Keane	Page 2 of 6

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <http://www.hse.ie/eng/Disclaimer>

This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens

Haematological:

Table 1: Recommended dose modification of vinorelbine based on neutrophil counts

ANC ($\times 10^9/L$)	Dose*
≥ 1.5	100% starting dose
1-1.49	50% starting dose
< 1	Delay one week and repeat FBC. If 3 consecutive weekly doses are held because neutrophil count is $< 1.0 \times 10^9/L$, discontinue vinorelbine
*Dose adjustment: For patients who during treatment, have experienced fever or sepsis while neutrophils were $< 1.5 \times 10^9/L$ or had 2 consecutive weekly doses held due to neutropenia, subsequent doses of vinorelbine should be given as follows	
ANC ($\times 10^9/L$)	Dose
≥ 1.5	75% starting dose
1-1.49	37.5% starting dose
< 1	Delay until count recovery and follow the above dosing guidelines

Renal and Hepatic Impairment:

Table 2: Dose modification of CISplatin and vinorelbine in renal and hepatic impairment

Drug	Renal Impairment		Hepatic Impairment		
CISplatin	Cr Cl (ml/min)	Dose	No dose reductions necessary		
	>60	100%			
	45-59	75%			
	<45	Delay			
Vinorelbine	No dose reduction necessary		AST/ALT	Bilirubin	Dose
			>5 x ULN	> 2 x ULN	Reduce dose by 33%
			ULN= Upper Limit of Normal		

Table 3: Dose modification schedule based on adverse events

Adverse reactions	Recommended dose modification
Peripheral neuropathy Grade 2	Withhold treatment until recovery to grade 1 then reduce the dose to 75% of the original dose.
Grade 3	Discontinue treatment
Grade 3 constipation	After appropriate management of symptoms (See supportive care) may consider reducing the dose of vinorelbine to 75% of the original dose.
Other toxicities \geq Grade 3	Defer therapy for 1 week until resolved to \leq grade 1. Discuss with consultant if > 1 week delay.

NCCP Regimen: Vinorelbine and CISplatin Therapy -28 day	Published: 20/06/2016 Review: 10/06/2025	Version number: 5
Tumour Group: Lung NCCP Regimen Code: 00343	ISMO Contributor: Prof Maccon Keane	Page 3 of 6

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <http://www.hse.ie/eng/Disclaimer>

This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens

SUPPORTIVE CARE:

EMETOGENIC POTENTIAL:

CISplatin High (**Refer to local policy**)
 Vinorelbine Minimal (**Refer to local policy**).

PREMEDICATIONS:

Pre and Post Hydration therapy required for CISplatin administration (**Reference local policy or see recommendations above**).

OTHER SUPPORTIVE CARE:

- Mouth care (**Refer to local policy**)
- Patients should be counseled on the risk of constipation associated with the use of vinca alkaloids. Dietary interventions or prophylactic laxatives may be required.
- Gastric protection with a proton pump inhibitor or a H₂ antagonist may be considered in patients considered at high risk of GI ulceration or bleed.

ADVERSE EFFECTS / REGIMEN SPECIFIC COMPLICATIONS

The adverse effects listed are not exhaustive. Please refer to the relevant Summary of Product Characteristics for full details.

- **Cardiac toxicity:** Special care should be taken when prescribing for patients with history of ischemic heart disease.
- **Extravasation:** Vinorelbine causes pain and tissue necrosis if extravasated (**Refer to local guidelines**).
- **Neutropenia:** The dose limiting adverse reaction of vinorelbine is mainly neutropenia. This effect is non-cumulative, having its nadir between 7 and 14 days after the administration and is rapidly reversible within 5 to 7 days. Fever or other evidence of infection must be assessed promptly and treated appropriately.
- **Constipation:** Constipation with vinorelbine should at a grade 1-2 be managed with dietary interventions or laxatives
- **Renal toxicity:** Renal toxicity is common with CISplatin. Encourage oral hydration. Avoid nephrotoxic drugs such as aminoglycoside antibiotics
- **Ototoxicity and sensory neural damage** should be assessed by history prior to each cycle.

DRUG INTERACTIONS:

- Risk of drug interactions causing increased concentrations of vinorelbine with CYP3A inhibitors.
- Risk of drug interactions causing decreased concentrations of vinorelbine with CYP3A inducers.
- CISplatin may potentiate the nephrotoxic and ototoxic effects of loop diuretics and aminoglycosides so concurrent use should be avoided.
- Current drug interaction databases should be consulted for more information

REFERENCES:

1. Winton T, Livingston R, Johnson D, et al. Vinorelbine plus cisplatin vs. observation in resected non-small-cell lung cancer. N Engl J Med 2005;352:2589-97.
2. BCCA Protocol Summary for Adjuvant CISplatin and Vinorelbine Following Resection of Non-Small Cell Lung Cancer LUJNP 1st April 2013
3. Nephrotoxicity Associated with CISplatin EviQ ID: 184 v.3

NCCP Regimen: Vinorelbine and CISplatin Therapy -28 day	Published: 20/06/2016 Review: 10/06/2025	Version number: 5
Tumour Group: Lung NCCP Regimen Code: 00343	ISMO Contributor: Prof Maccon Keane	Page 4 of 6

The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at <http://www.hse.ie/eng/Disclaimer>

This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens

<https://www.eviq.org.au/clinical-resources/side-effect-and-toxicity-management/prophylaxis-and-prevention/184-nephrotoxicity-associated-with-CISplatin>

4. Portilla D et al. CISplatin nephrotoxicity. UpToDate Accessed Oct 2017
https://www.uptodate.com/contents/CISplatin-nephrotoxicity?source=search_result&search=CISplatin%20hydration&selectedTitle=1~150
5. Dosage Adjustment for Cytotoxics in Renal Impairment January 2009; North London Cancer Network.
6. Dosage Adjustment for Cytotoxics in Hepatic Impairment January 2009; North London Cancer Network.
7. Krens S D, Lassche, Jansman G F G A, et al. Dose recommendations for anticancer drugs in patients with renal or hepatic impairment. Lancet Onco/2019; 20:e201-08. [https://doi.org/10.1016/S1470-2045\(19\)30145-7](https://doi.org/10.1016/S1470-2045(19)30145-7)
8. Cisplatin 1mg/ml Concentrate for Solution for Infusion. Summary of Product Characteristics Last updated: 11/03/2019. Accessed June 2020. Available at <https://www.hpra.ie/img/uploaded/swedocuments/Final%20approved%20SPC%20PA0822.199.001.pdf>
9. Vinorelbine (Navelbine[®]) Summary of Product Characteristics. Last updated: 24/09/2019. Accessed Jun 2020 Available at https://www.hpra.ie/img/uploaded/swedocuments/Licence_PA0329-011-003_24092019091922.pdf
10. NCCP Classification Document for Systemic Anti-Cancer Therapy (SACT) Induced Nausea and Vomiting. V3 2021. Available at: <https://www.hse.ie/eng/services/list/5/cancer/profinfo/chemoprotocols/nccp-classification-document-for-systemic-anti-cancer-therapy-sact-induced-nausea-and-vomiting.pdf>
11. Adjuvant vinorelbine plus cisplatin for resected early stage non-small cell lung cancer. UpToDate: Version 23.0. Available at: https://www.uptodate.com/contents/image?imageKey=ONC%2F59705&topicKey=ONC%2F85689&search=Vinorelbine%20cisplatin%20lung&source=outline_link&selectedTitle=2~150 Accessed Jun 2020.

Version	Date	Amendment	Approved By
1	20/06/2016		Prof Maccon Keane
2	20/06/2018	Updated with new NCCP regimen template. Update of treatment table to: <ul style="list-style-type: none"> • Standardise fluid and infusion times. • Amend CISplatin hydration recommendations Update to dosing in hepatic and renal impairment Emetogenic potential updated	Prof Maccon Keane
3	21/08/2019	Updated recommended dose modification of vinorelbine based on neutrophil counts	Prof Maccon Keane
4	10/06/2020	Reviewed.	Prof Maccon Keane
5	24/06/2020	Updated CISplatin hydration protocol	Prof Maccon Keane

Comments and feedback welcome at oncologydrugs@cancercontrol.ie.

NCCP Regimen: Vinorelbine and CISplatin Therapy -28 day	Published: 20/06/2016 Review: 10/06/2025	Version number: 5
Tumour Group: Lung NCCP Regimen Code: 00343	ISMO Contributor: Prof Maccon Keane	Page 5 of 6
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		

NCCP Regimen: Vinorelbine and Cisplatin Therapy -28 day	Published: 20/06/2016 Review: 10/06/2025	Version number: 5
Tumour Group: Lung NCCP Regimen Code: 00343	ISMO Contributor: Prof Maccon Keane	Page 6 of 6
<p>The information contained in this document is a statement of consensus of NCCP and ISMO or IHS professionals regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult these documents is expected to use independent medical judgement in the context of individual clinical circumstances to determine any patient's care or treatment. Use of these documents is the responsibility of the prescribing clinician and is subject to HSE's terms of use available at http://www.hse.ie/eng/Disclaimer</p> <p><i>This information is valid only on the day of printing, for any updates please check www.hse.ie/NCCPchemoregimens</i></p>		